

References

1. Allander T, Tammi MT, Eriksson M, Bjerkner A, Tiveljung-Lindell A, Andersson B. Cloning of a human parvovirus by molecular screening of respiratory tract samples. *Proc Natl Acad Sci U S A*. 2005;102:12891–6.
2. Sloots TP, McErlean P, Speicher DJ, Arden KE, Nissen MD, Mackay IM. Evidence of human coronavirus HKU1 and human bocavirus in Australian children. *J Clin Virol*. 2006;35:99–102.
3. Ma X, Endo R, Ishiguro N, Ebihara T, Ishiko H, Ariga T, et al. Detection of human bocavirus in Japanese children with lower respiratory tract infections. *J Clin Microbiol*. 2006;44:1132–4.
4. Bastien N, Brandt K, Dust K, Ward D, Li Y. Human bocavirus infection, Canada. *Emerg Infect Dis*. 2006;12:848–50.
5. Foulongne V, Rodiere M, Segondy M. Human bocavirus in children. *Emerg Infect Dis*. 2006;12:862–3.
6. Arnold JC, Singh KK, Spector SA, Sawyer MH. Human bocavirus: prevalence and clinical spectrum at a children's hospital. *Clin Infect Dis*. 2006;43:283–8.
7. Weissbrich B, Neske F, Schubert J, Tollmann F, Blath K, Blessing K, et al. Frequent detection of bocavirus DNA in German children with respiratory tract infections. *BMC Infect Dis*. 2006;6:109.
8. Chung JY, Han TH, Kim CK, Kim SW. Bocavirus infection in hospitalized children, South Korea. *Emerg Infect Dis*. 2006;12:1254–6.
9. Choi EH, Lee HJ, Kim SJ, Eun BW, Kim NH, Lee JA, et al. The association of newly identified respiratory viruses with lower respiratory tract infections in Korean children, 2000–2005. *Clin Infect Dis*. 2006;43:585–92.

Address for correspondence: Marjolein Y. Berger, Erasmus MC, Department of General Practice, Room Ff 332, PO Box 2040, 3000 CA Rotterdam, the Netherlands; email: m.berger@erasmus.nl

Dengue Virus Serotype 3, Karachi, Pakistan

To the Editor: The global prevalence of dengue fever (DF) has grown dramatically in recent decades; DF is now endemic to >100 countries (*1*). Dengue hemorrhagic fever (DHF), a potentially lethal complication of dengue virus infection, was first recognized in Asia in the 1950s and is now a leading cause of hospitalization and death among children (*1*). During the past decade, DHF epidemics have occurred in China, Sri Lanka, India, the Maldives, Bangladesh, and Pakistan (*2–4*).

In Pakistan, an outbreak of DHF was first reported in Karachi in 1994 (*4*). Through mid-2005, 15–20 patients with DF or DHF were admitted each year to the Aga Khan University Hospital (AKUH), a tertiary care referral center in Karachi. Many more cases, however, may have gone unrecognized. Ours is the first report of dengue virus serotype 3 in Pakistan.

From September through December 2005, at least 3 major hospitals in Karachi, including AKUH, had a sudden increase in the number of patients with signs consistent with the World Health Organization definition of DHF: high fever, rash, epistaxis, gum bleeding, liver dysfunction, and thrombocytopenia (platelets <100,000/mm³); most had evidence of capillary leakage in the form of raised hematocrit and pleural effusion with or without ascites (*5*). Because in Pakistan, Crimean-Congo hemorrhagic fever (CCHF) is an important differential diagnosis for hemorrhagic fever, most patients seen at AKUH received care in strict isolation and were empirically treated with ribavirin. At time of admission, blood samples were collected for serologic testing for dengue virus and reverse transcription (RT)–PCR testing for

CCHF virus. The first 5 samples, collected during the initial 2 weeks of the outbreak, were also sent to the Special Pathogens Reference Unit, Centre for Emergency Preparedness and Response, Health Protection Agency, Salisbury, United Kingdom, for diagnostic confirmation. In the absence of a local surveillance and disease notification system, the number of patients with suspected DHF at different hospitals in Karachi could not be ascertained.

Of the 106 patients who had a clinical diagnosis compatible with DHF (*5*), 9 (8.5%) died and 97 (91.5%) recovered. Patients with possible DF (fever, mild thrombocytopenia with platelets >100,000/mm³) were not admitted and were treated as outpatients. Dengue virus infection was confirmed for 42 of the 106 patients. Serum samples from 39 patients contained anti-dengue virus immunoglobulin M (IgM) antibody (Chemicon, Temecula, CA, USA). Diagnosis for 6 of these patients was confirmed by using immunoblot tests (Dengue IgM Blot and Dengue IgG Blot, Genelabs Diagnostics, Singapore). Of the 9 patients who died, 6 had dengue IgM and IgG according to immunoblot testing, and 3 had dengue IgM according to ELISA. Diagnoses for 3 additional patients were confirmed by RT-PCR.

An RT-PCR assay specific for dengue viruses (*6*) was used to amplify the C/PrM/M region of the genome and produced PCR products of the expected size in 3 patient samples: 2 (K1 and 2) from Karachi and 1 (B) from Balochistan. The PCR products were sequenced, and data were subsequently placed in GenBank under accession numbers DQ469827 for D3418-05 (patient K1), DQ469828 for D3419-05 (patient K2), and DQ469826 for D3417-05 (patient B). These data were compared with those in databases by using the basic local alignment search tool for nucleotides (blastn), with default settings (*7*). For

EID
Online
www.cdc.gov/eid

each sequence analyzed, the lowest Expect (E) value showed significant similarity with a dengue serotype 3 isolate from India in 2004 (DQ323042). A phylogenetic tree was constructed with a collection of dengue sequences (online Appendix Table, available from www.cdc.gov/ncidod/EID/13/1/182-appT.htm). Phylogenetic relationships between sequences are depicted in the Figure. Sequences from the 2005 outbreak are most similar to those from Indian strains of dengue serotype 3, which were isolated in Delhi.

An unexpected finding was the detection, at both AKUH and the UK Special Pathogens Reference Unit, of dengue-3 and CCHF virus RNA in the sample from patient B. CCHF is endemic to the rural Balochistan province of Pakistan, where DF has

been documented (8). In the absence of information on the current dengue situation in Balochistan and given the increasing dengue activity in Karachi, a similar increase can be assumed for Balochistan. Possible introduction of dengue serotype 3 in a CCHF-endemic area resulted in dual infection in patient B, who essentially had clinical and laboratory features compatible with DHF. Patient B received ribavirin and recovered. Our results suggest that the 2005 outbreak of DHF in Karachi, Pakistan, was caused by strains of dengue virus serotype 3 related to those circulating in India (9).

Acknowledgments

We thank Aslam Khan, Maqsood Bhatti, and Roshan Hadwani for their contributions to this work.



Figure. Maximum-likelihood phylogenetic tree of established dengue virus 3 serotypes and new sequences from Pakistan identified in this study. The tree is based on a 238-nt sequence alignment comprising the C/PrM/M gene (nucleotides: 179–417 dengue virus 3 prototype [NC_001475]). The bar shows the number of substitutions per bases weighted by the Tamura-Nei algorithm. Horizontal distances are equivalent to the distances between isolates; numbers at nodes indicate support values for the branch of the tree inferred at the node. The origins of the sequences used to reconstruct this tree are detailed in the online Appendix Table.

**Bushra Jamil,* Rumina Hasan,*
Afia Zafar,* Kevin Bewley,†
John Chamberlain,† Valerie
Mioulet,† Moira Rowlands,†
and Roger Hewson†**

*Aga Khan University Hospital, Karachi, Pakistan; and †Health Protection Agency, Salisbury, United Kingdom

References

1. World Health Organization. Dengue and dengue haemorrhagic fever. Fact sheet no. 117; Revised 2002 Apr [cited 2006 Oct 31]. Available from <http://www.who.int/mediacentre/factsheets/fs117/en/>
2. Kabilan L, Balasubramanian S, Keshava SM, Satyanarayana K. The 2001 dengue epidemic in Chennai. *Indian J Pediatr*. 2005;72:919–23.
3. Islam MA, Ahmed MU, Begum N, Choudhury NA, Khan AH, Parquet MC, et al. Molecular characterization and clinical evaluation of dengue outbreak in 2002 in Bangladesh. *Jpn J Infect Dis*. 2006;59:85–91.
4. Chan YC, Salahuddin NI, Khan J, Tan HC, Seah CL, Li J, et al. Dengue haemorrhagic fever outbreak in Karachi, Pakistan, 1994. *Trans R Soc Trop Med Hyg*. 1995;89:619–20.
5. Dengue haemorrhagic fever: diagnosis, treatment, prevention and control. 2nd ed. Geneva: World Health Organization, 1997.
6. Lanciotti RS, Calisher CH, Gubler DJ, Chang G-J, Vorndam V. Rapid detection and typing of dengue viruses from clinical samples using reverse transcriptase-polymerase chain reaction. *J Clin Microbiol*. 1992;30:545–51.
7. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic Local Alignment Search Tool. *J Mol Biol*. 1990; 215:403–10.
8. Paul RE, Patel AY, Mirza S, Fisher-Hoch SP, Luby SP. Expansion of epidemic dengue viral infections to Pakistan. *Int J Infect Dis*. 1998;2:197–201.
9. Akram DS, Igarashi A, Takasu T. Dengue virus infection among children with undifferentiated fever in Karachi. *Indian J Pediatr*. 1998;65:735–40.

Address for correspondence: Bushra Jamil, Assistant Professor, Department of Medicine, Pathology, and Microbiology, Aga Khan University Hospital, Stadium Rd, PO Box 3500, Karachi 74800, Pakistan; email: jamil.bushra@aku.edu